

Inertia, coarsening and fluid motion in binary mixtures

M. E. Cates^a, V. M. Kendon^a, P. Bladon^a, J.-C. Desplat^b

^a*Department of Physics and Astronomy and* ^b*Edinburgh Parallel Computing Centre,
University of Edinburgh, King's Buildings, Mayfield Road, Edinburgh EH9 3JZ, Scotland*

Symmetric binary fluids, quenched into a regime of immiscibility, undergo phase separation by spinodal decomposition. In the late stages, the fluids are separated by sharply defined, but curved, interfaces: the resulting Laplace pressure drives fluid flow. Scaling ideas (of Siggia and of Furukawa) predict that, ultimately, this flow should become turbulent as inertial effects dominate over viscous ones. The physics here is complex: mesoscale simulation methods (such as Lattice Boltzmann and Dissipative Particle Dynamics) can play an essential role in its elucidation, as we describe. Likewise, it is a matter of experience that immiscible fluids will mix, on some lengthscale at least, if stirred vigorously enough. A scaling theory (of Doi and Ohta) predicts the dependence of a steady state domain size on shear rate, but assumes low Reynolds number (inertia is neglected). Our preliminary simulation results (three-dimensional, so far only on small systems) show little sign of the kind of steady state envisaged by Doi and Ohta; they raise instead the possibility of an oriented domain texture which can continue to coarsen until either inertial effects, or (in our simulations) finite size effects, come into play.

I. INTRODUCTION

When an incompressible binary fluid mixture is quenched far below its spinodal temperature, it will phase separate into domains of the two fluids. The simplest case (in theory at least) is when the two fluids are thermodynamically and kinetically symmetrical: fluids A and B have identical properties in all respects except that they are mutually phobic. For a deep quench (to a temperature well below the critical point) the thermodynamic equilibrium state then has the two fluids completely demixed. One remaining control parameter is the volume fraction ϕ ; only for a 50:50 mix ($\phi = 0.5$, which with thermodynamic symmetry is a quench through the critical point) is the system totally symmetrical. Other significant parameters are the surface tension σ , viscosity η and the fluid mass density ρ . There is also a mobility parameter M which controls the collective diffusion of concentration fluctuations: this can remain important at late times, but only if ϕ is small enough (less than about 0.15–0.20 for 3-D) that the fluid domains depercolate. The minority droplets can then continue to coarsen by a ripening mechanism (controlled by M) or by diffusion and coalescence of the droplets themselves; both give a mean droplet size that scales with time as $t^{1/3}$. If instead the domains remain connected, then the late-stage growth is driven by capillary forces (Laplace pressure), arising from curvature of the (sharply defined) interface between the two fluids; these drive fluid flows from regions of tight curvature (necks, narrow liquid bridges) into those of low interfacial curvature (large domains). Note that for coarsening of a bicontinuous structure to proceed this way, one also requires discrete “pinch-off” events to continually occur: each of these allows the topology to change discontinuously in time. For a theory-oriented review of the late stages of spinodal domain growth, see Bray [1]; there are many relevant experimental studies as well [2].

In what follows, we address two issues concerning domain growth in three dimensional binary fluid systems. (The two dimensional case has some special features of its own; see [3].) The first issue concerns the role of inertia in the late stages; this is a matter of continuing interest and controversy [4], that our recent simulation work [5] illuminates. The second concerns the effect of an applied shear flow on the coarsening process. Here we have only preliminary simulation results, but these are enough to suggest that there is more to this problem than a recent scaling theory [6] suggests; again, our discussion focuses on inertial effects.

Our simulation work uses two different mesoscale simulation methods. One is the Lattice Boltzmann (LB) method, in which a velocity distribution function $f(\mathbf{v})$ and composition variable ϕ are defined at discrete lattice points. The rules for updating these quantities represent a discretization of the (zero temperature) Navier-Stokes equation for fluid flow, coupled to a Cahn-Hilliard type thermodynamic free energy functional $\mathcal{F}[\phi]$, which provides the driving force for phase separation. The second method is DPD (dissipative particle dynamics), which is a long-time-step, noisy molecular dynamics algorithm (off lattice) involving soft repulsive interaction potentials between two types of particle. (The repulsive interactions are chosen to favour demixing.) This is a fixed-temperature algorithm, in which stability is ensured, despite the long time-step, by introducing local damping and noise terms in accord with a suitable fluctuation dissipation theorem. Crucially, both damping and noise act pairwise on particle velocities and hence conserve momentum locally, recovering the isothermal Navier Stokes equation at large length scales. For further information on LB for binary fluids see Swift et al. [7] and on DPD see Groot and Warren [8]; details of our algorithms and parameter settings are recorded elsewhere [5,9,10]. Note that in both methods we have run tests to check that,

under the particular simulation conditions we adopt, fluids are behaving incompressibly.

II. BINARY DEMIXING AT HIGH REYNOLDS NUMBER

For simplicity, we restrict attention to fully symmetric mixtures ($\phi = 0.5$) for which the fluid domains comprise, in three dimensions, a fully bicontinuous structure. The late-time evolution of this structure remains incompletely understood despite theoretical [11,12,1,4], experimental [2] and simulation [13,14,10,15] work over recent years.

A. Scaling expectations

As emphasized by Siggia [11], the coarsening of a bicontinuous spinodal texture involves capillary forces (governed by the interfacial tension, σ), viscous dissipation (governed by the fluid viscosity η), and fluid inertia (governed by the mass density ρ). Out of these three physical parameters, (σ , η , and ρ), only one length, $L_0 = \eta^2/\rho\sigma$ and one time $T_0 = \eta^3/\rho\sigma^2$ can be constructed, which allow us to describe the time evolution of the coarsening system in unique dimensionless length and time measures. For convenience we define the lengthscale $L(T)$ of the domain structure at time T via the structure factor $S(k)$ as $L = (1/2\pi) \int S(k)dk / \int kS(k)dk$. Provided no other physics except that described by the three macroscopic parameters, σ , η , and ρ , is involved in late stage growth this leads us to the dynamical scaling hypothesis [11,12]:

$$l = l(t) \tag{1}$$

where we use reduced time and length variables, $l \equiv L/L_0$ and $t \equiv (T - T_{int})/T_0$. Since dynamical scaling should hold only after interfaces have become sharp, and transport by molecular diffusion ignorable, we have allowed for a nonuniversal offset T_{int} ; thereafter the scaling function $l(t)$ should, in principle, approach a universal form, the same for all (fully symmetric, deep-quenched, incompressible) binary fluid mixtures.

It was argued further by Furukawa [12] that, for small enough t , fluid inertia is negligible compared to viscosity, whereas for large enough t the reverse is true. Dimensional analysis then requires the following asymptotes:

$$l \rightarrow bt \quad ; \quad t \ll t^* \tag{2}$$

$$l \rightarrow ct^{2/3} \quad ; \quad t \gg t^* \tag{3}$$

where, if dynamical scaling holds, amplitudes b, c and the crossover time t^* (defined, for example, by the intersection of asymptotes on a log-log plot) are universal.

Several earlier numerical studies claim to see one or other of these scaling regimes, but of these, few provide accurate values of η, ρ, σ as required if the various datasets are to be compared on a scaling plot [15]. Those that do so, and which appear to confirm Eq.2, include the work of Laradji et al. [13], for which $2 \leq l \leq 20$, that of Bastea and Lebowitz [14] ($1 \leq l \leq 2$) and that of Jury, Bladon, Krishna and Cates [10] ($20 \leq l \leq 2000$). Jury et al. performed a careful comparison of their own (DPD) and these others' datasets; while linear scaling was reported, they found no consistency in the value of b . Instead, a systematic trend appeared, in which b drifts downward between datasets as one moves to larger t and l (roughly, $b \sim t^{-0.2}$). Despite this, within each DPD dataset, the linear t dependence is clearly better than a fit to $l \sim t^{0.8}$. Jury et al. proposed that a non-scaling behaviour of this kind could perhaps be explained if some nonuniversal physics (that is, not contained in σ, ρ, η) were to intervene: they suggested a candidate involving the physics of topological reconnection, a process that, even at late times, could involve a molecular (or discretization) scale, small compared to L .

This possibility remains open, although our own, more recent, LB data [5] suggests the following alternative explanation: (a) the linear law is indeed obeyed for $l \leq 20$ (with $b \simeq 0.07$), but the b coefficients of Laradji et al. and of Bastea and Lebowitz are both overestimated due to residual diffusion effects; (b) the data of Jury et al. lies within a broad crossover region between Eq.2 and Eq.3, where the local slope on a log-log plot is around 0.8; (c) the preference for linear fits in these DPD datasets is partly caused by finite size corrections within each dataset. To better eliminate the latter, in our LB coarsening data we insist that $L \leq \Lambda/4$, with Λ the linear system size (Jury et al. allowed $L \leq \Lambda/2$). A comparison between representative LB and DPD datasets, in a regime where both are available, is shown in Fig. 1. Though not identical, it is hard to be sure that the remaining discrepancies do not arise from finite size corrections.




FIG. 1. Comparison of individual $l(t)$ datasets, on a log-log plot, generated by the DPD [10] and the LB [5] algorithms. LB dataset (crosses) has $L_0 = 0.15$ lattice units, DPD datasets (solid) have $L_0 = 0.29, 0.19, 0.13, 0.077$ DPD units (left to right).

B. Inertial effects

The relative importance of inertial to viscous terms, in fluid mechanical problems, is traditionally measured by a Reynolds number. In the spinodal context this is usually defined as $Re = Re_s \equiv (\rho L / \eta) dL/dT = \dot{l}l$. In the linear regime it is of order $b^2 t$. Note that since b is rather small, the maximum Re achieved by Jury et al (for $t \simeq 2000$) is only about 20 even with t is of order $10^4 - 10^5$. Indeed, this is roughly where we now believe t^* to lie; a Reynolds number of 20 is large enough for inertial effects to be non-negligible, but by fluid mechanics standards, still modest. Note also that the scaling ideas clearly predict that Re should increase without bound (as $t^{1/3}$) within the inertial regime, Eq.3. However, in a recent paper [4] Grant and Elder have argued that the Reynolds number cannot, in fact, continue to grow indefinitely. If so, Eq.3 is not truly the large t asymptote, which must instead have $l \sim t^\alpha$ with $\alpha \leq \frac{1}{2}$. Essentially, Grant and Elder argue that at large enough Re , turbulent remixing of the interface will limit the coarsening rate [4], so that Re stays bounded at a level which they estimate as $Re \sim 10 - 100$.

Our recent LB work [5] represents the first large-scale simulations of 3-D spinodal decomposition to unambiguously attain a regime in which inertial forces dominate over viscous ones, allowing a test of this idea. We find direct evidence for Furukawa's $l \sim t^{2/3}$ scaling, Eq.3. Although a further crossover to a regime of saturating Re cannot be ruled out, we find no evidence for this up to $Re \simeq 350$. Two of our LB datasets are shown in Fig.2. The first has high L_0 (high viscosity) and corresponds to l values around unity ($t \simeq 10$) for the fitted part of the data. It is well fit by Eq.2. The second has low L_0 and corresponds to l around 10^5 (or $t \simeq 10^7$); it is well fit by Eq.3. The Reynolds number is about 0.1 near the end of the first run and about 350 near the end of the second.

It goes without saying that no *single* simulation run could possibly cover this range of length or time scales, since that would require a lattice of around $(10^6)^3$ sites. Our lattices are 256^3 , but by changing the LB parameter values, we can access both regimes, and everything in between [5]. Similar exploration is possible (to a lesser extent) within DPD [10]. Such mesoscale methods have strong advantages over (say) MD [13] since, after proper calibration and subject to specified range limitations, they allow one to “dial in” one's own choice of thermodynamic (σ) and kinetic (η) parameters. Thus one can build up the $l(t)$ curve section by section; if it is universal, this is enough. (For a fuller explanation of the procedure, see Kendon et al. [5].) This is done for our LB data in Fig.3; note that the rightmost two datasets (both in the inertial regime) are almost contiguous, as one would hope on a universal scaling plot. (Each has a slope close to $2/3$.) However, computational restraints prevent us so far from covering the entire $l(t)$ curve with data; it remains possible that some mismatches between runs, of the type reported by Jury et al. [10] might still be observed at intermediate l, t . Note the broad crossover between viscous (three leftmost runs) and inertial




FIG. 2. L vs. T (in lattice units) for runs with $L_0 = 5.9$ (circles) and 0.0003 (diamonds). Dotted lines delineate the range of data points used for fitting ($17 < L < 64$) and the fits to t^1 and $t^{2/3}$ respectively have been projected back to show the intercepts, T_{int} .

(two rightmost runs) regimes: this crossover covers four decades in t (or three in l). But it is less spectacular when expressed in terms of Re ; our data span $0.1 \leq Re \leq 350$ and the crossover region is roughly $Re = 1 - 100$.




FIG. 3. Scaling plot in reduced variables (l, t) for eight LB datasets. Second and third from left overlap and cannot be distinguished on the plot.

It is important now to ask whether, at the largest Re values we can reach, there is significant turbulence in the fluid flow. One quantitative signature of turbulence is the skewness S of the longitudinal velocity derivatives; this is close to zero in laminar flow but approaches $S = -0.5$ in fully developed turbulence [16]. A space-averaged value close to $S = -0.5$ would imply turbulence everywhere and, presumably, significant remixing at the interface [4]. We do detect increasingly negative S as Re is increased but reach only $S \simeq -0.35$ for $Re \simeq 350$ [17]. This suggests that at our highest Re 's, partially but not fully developed turbulence is present. This view is consistent with Fig.4, which shows velocity maps for low and high Re runs, and (for comparison) single fluid freely decaying isotropic turbulence. In a low Re flow one expects the only relevant lengthscale in the system to be the domain size L ; at high Re , there should

be a cascade of structure in the velocity field below the domain size. Some internal structure is plausibly, though not conclusively, visible in the plot for our lowest L_0 value. But since we do not yet see fully developed turbulence, it remains an open issue whether such effects would lead, beyond the observed $t^{2/3}$ inertial scaling regime, to a final, turbulent remixing regime (of saturating Re) as proposed by Grant and Elder [4]. If it does, the limiting value of Re must significantly exceed their estimate of $10 - 100$.




FIG. 4. Velocity maps for $L_0 = 5.9, 3.0 \times 10^{-4}$ (LB), and for freely decaying isotropic turbulence (pseudo-spectral dns). Each map projects a $32 \times 32 \times 2$ thin section taken from a 128^3 grid. The interface appears wide where it cuts the section at a glancing angle; each site shows two arrows which are the projected velocity vectors in the two layers of the section.

III. DEMIXING UNDER FLOW

Consider now the situation where a symmetric binary fluid undergoes a deep quench in conditions where a steady shear flow is applied externally. In simulations, there are two ways to achieve this. One is by Lees-Edwards (moving periodic) boundary conditions, which have been implemented in our DPD codes [18]; the other is to have sliding solid boundaries, and the LB results reported below use this method. For binary fluids under shear, the two approaches give qualitatively similar results. For related work in two and three dimensions, see [20,21]. The questions are as follows: does the presence of flow arrest the coarsening process, giving a steady state with a finite domain size, and, if so, what is the character of this steady state? Also, does this steady state depend on starting conditions – for example, if the binary fluids are allowed to demix first and the flow is then switched on, is the same final state achieved? Perhaps surprisingly (despite the obvious importance of mixing and emulsification in many technological areas) well-founded theoretical answers to these questions are almost nonexistent.

A. The Doi-Ohta theory

An important theoretical inroad into this problem was made by Doi and Ohta [6]. (Its rheological consequences are reviewed in the book by Larson [19], where comparisons to experiment can be found.) The Doi-Ohta work is entirely limited, however, to a regime of vanishingly small Reynolds number. Note that for this problem, the relevant Re is the larger of $\text{Re}_s = (\rho L / \eta) dL / dT$ (the value extracted from the coarsening process, which must vanish in any true steady state) and

$$\text{Re}_f = \rho \dot{\gamma} L^2 / \eta \quad (4)$$

which is the value found by considering the effect of the external shear rate $\dot{\gamma}$ on a droplet of scale L . Both definitions assume, perhaps implausibly, that there is only one relevant length, L , which requires, for example, that domains are not strongly anisotropic; we return to this point below.

Doi and Ohta assumed that, for nonzero $\dot{\gamma}$, a steady state would be reached at negligible Re (so that inertial effects can be ignored). The applied flow stretches and deforms the interfaces (tending also to orient these along the flow direction); this is opposed by interfacial tension. The competition is measured by the capillary number $Ca = T_0 \dot{\gamma} L / L_0 = \dot{\gamma} L \eta / \sigma$; at small Ca , a spherical droplet of size L is weakly perturbed, but at large Ca it should be elongated to the point of rupture. By estimating the magnitude of each effect in the absence of the other, and balancing these, Doi and Ohta made the intriguing prediction that, in steady state, the fluid should have a $\dot{\gamma}$ -independent viscosity of the form $\eta f(\phi)$. That is, the behaviour should be independent of σ and ρ , but dependent on ϕ (values of ϕ close to 0.5 were assumed.) This form follows purely by dimensional analysis if one insists that the only relevant variables are $\dot{\gamma}, \sigma, \eta$ and ϕ (the absence of ρ from this list is because inertia is neglected.) Doi and Ohta went on to develop specific approximation schemes that allowed them, for example, to study transient (shear startup) behaviour.

In the steady state, the departure of $f(\phi)$ from unity must come from the presence of interfacial stresses, which, if the structure is characterized by a single domain length L , scale as σ/L . Comparing this with the excess shear stress $\eta \dot{\gamma}(f - 1)$ we obtain a length estimate in steady state

$$L \simeq \frac{\sigma}{\eta \dot{\gamma}} \quad (5)$$

which corresponds to the system arranging itself, in steady state, so that the capillary number is of order unity. This argument is, of course, somewhat oversimplified. If more than one length scale is present (for example, describing the structure along different directions relative to the flow) then one has to determine which one dominates the interfacial contribution to the viscous stress. The largest Laplace stresses arise in regions of strongest curvature; however, if the strongest curvature is axial about the flow direction (for example, thin cylinders lying along the flow) it cannot contribute to the shear stress at all [6]. Doi and Ohta did not much explore these geometrical aspects of their work. Note that, implicit in their whole description, is an assumption that a steady state can be achieved without inertial terms coming into play. Our preliminary results on sheared spinodal systems, given below, cast some doubt on this assumption.

B. Results

We have simulated sheared binary fluids using both DPD and LB algorithms. The results presented here are for relatively small systems (50,000 DPD particles, and 64^3 or 96^3 lattices) and thus are preliminary, but we think they point towards some interesting physics. In future work we hope to address larger systems; among other things, this will allow reliable statistics for the interfacial stresses, and hence for the rheological properties of the sheared systems, to be obtained.

Our first result is that shear does inhibit coarsening. If one defines a single lengthscale via the total surface area A in the system (so that $A = \Lambda^3/L$), then indeed this lengthscale increases more slowly, when shear is applied, than without it (Fig.5).

On the other hand, this one-lengthscale analysis fails to reflect the extreme anisotropy that can develop. Indeed, by examining the eigenvalues $\lambda_1, \lambda_2, \lambda_3$, of the interfacial curvature matrix [21],

$$D_{\alpha\beta} = \frac{\sum_{grid} \partial_\alpha \hat{\phi} \partial_\beta \hat{\phi}}{\sum_{grid} \hat{\phi}^2}, \quad (6)$$

lengths for the three different directions can be estimated,

$$L_1 = \frac{\xi}{\lambda_1}, \quad L_2 = \frac{\xi}{\lambda_2}, \quad L_3 = \frac{\xi}{\lambda_3}. \quad (7)$$

In these expressions, ξ is the equilibrium width of the fluid-fluid interface, and $\hat{\phi}$ is the local compositional order parameter defined on the lattice. The time evolutions of these three lengths are plotted for a system with $\dot{\gamma} T_0 = 0.14$ in Fig.6.

This data shows strong anisotropy developing, with elongation of domains along the flow direction. Under these conditions, finite size effects become greatly emphasized [22]: domains, elongated along the flow, soon reach the size of the simulation box and connect up to their periodic images. A sheared LB configuration, during the initial phase




FIG. 5. Overall coarsening of the domains as estimated from the amount of interface in an LB system for shear rates of $\dot{\gamma}T_0 = 0.07, 0.14, 0.21$, compared to zero shear. Using the system size to estimate the maximum capillary number for the systems due to the shear gives $Ca = 0.4, 0.8, 1.5$.




FIG. 6. Time evolution of three length scales given by the eigenvalues of the curvature matrix for the LB system with $\dot{\gamma}T_0 = 0.14$.

before this happens (such that $L_{1,2,3}$ are all small compared to the simulation size Λ) is shown in Fig.7(a). However, this configuration is not close to a steady state.

The structure we observe on approach to the steady state is, in all cases we have so far studied, clearly finite-size influenced. For ϕ different from 0.5, the appearance is like that of a “string phase” [23] in colloids: effectively infinite tubes of the minority fluid oriented along the streamlines. A typical example of this, found using DPD, is shown in Fig.7(b). Although no longer bicontinuous in the transverse direction, this structure can continue to coarsen, very slowly, by diffusive transport (ripening): it seems likely that, for this system size, the final state will be a single

cylinder of fluid. Figure 7(c) shows a LB simulation for $\phi = 0.5$ viewed along the streamlines; this has coarsened to the point where finite size effects are obvious even transverse to the flow direction. The structure is strongly oriented, with continuous domains of the two fluids passing right through the sample from front to back; however, it remains connected transverse to the flow with thin necklike bridges clearly visible. In fact, for $\phi = 0.5$, such transverse connectivity is inevitable unless the symmetry between the two fluids is spontaneously broken. Thus the structure remains capable of coarsening further, by a non-diffusive hydrodynamic mechanism. For this system size, the final state could be total separation, just as it would be without shear flow: such a state might be stable so long as the final interface between the two fluids is parallel to the streamlines.




FIG. 7. (a) LB simulation with $\phi = 0.5$, prior to steady state being achieved. Shown is a central 32^3 section from a 64^3 system, with the shear applied through the top plane moving left and the bottom plane moving right. Light blue denotes the fluid-fluid interface. The horizontal plane shows a slice through the order parameter profile, with solid colours representing the two fluids: this shows the interfacial sharpness. (b) DPD simulation with $\phi = 0.2$ and $\dot{\gamma}T_0 = 0.32$. Tubes are oriented along the streamlines. (c) LB simulation with $\phi = 0.5$ and $\dot{\gamma}T_0 = 0.14$. This view is along the streamlines; the top and bottom edges are the moving boundaries. The block colours (as in (a)) show the identity of the two fluids, now on a vertical plane towards the back of the simulation cell. Passing in front of this, one observes narrow fluid necks connecting blocks of similar fluid.

C. Discussion

In our simulations on these (fairly small) systems, we see no evidence of a steady state structure emerging that is independent of system size. This is intriguing. If the Doi-Ohta scaling is correct, we should obtain a length scale given by Eq.5; so long as this is small compared to Λ (and the steady state structure is not too anisotropic) the simulation should achieve a proper steady state, representative of that of an infinite sample. Put differently, this will arise if the maximum capillary number attainable in a system of size Λ , $Ca = \dot{\gamma}\Lambda\eta/\sigma$, is much larger than one. This is true for our largest DPD runs, with $Ca \sim 7$, yet we see no sign of any steady state not dominated by finite size effects; larger runs would be useful to clarify this issue.

An important possibility, which these results suggest, is as follows. By neglecting inertia and yet assuming a steady state, Doi and Ohta implicitly assume that this steady-state can be characterized by a notional capillary number $Ca = \dot{\gamma}L\eta/\sigma$ (based on a single measure of domain size L) of order one. This might be incorrect: without inertia, there may simply be no obstacle to complete separation of the two fluids in simple shear. Indeed, one can instead postulate a fully phase separated state with two blocks of fluid, separated by an interface in the plane of shear (or, in fact, any other orientation with the surface normals everywhere perpendicular to the fluid velocity). For such a flow, the *notional* capillary number is infinite (since L is infinite), but the *effective* value is zero: the interface is oriented parallel to the streamlines and there is no stretching of it by the flow. This structure is likely to be destabilized by inertial contributions, since it involves steady shear of a laminar interface between two fluids (although this may require a threshold of shear rate, such as $\dot{\gamma}T_0 \simeq 1$, to be exceeded). Thus it remains possible that a steady state, with a non-infinite domain scale can arise *only* when inertial terms are significant. If so the Reynolds number is just as important as the capillary number, and the implicit criterion $Ca \simeq 1$ of the Doi-Ohta theory is not adequate. To

test this point, one needs to run simulations under shear at high Re. We intend to pursue this, using our parallel LB code [9], in the near future.

Meanwhile it is straightforward to check that, at the system sizes and shear rates used here, a starting configuration with oriented slabs of fluid (with an interface parallel to the shear planes) remains essentially unperturbed by the flow. However, if the initial interface normal is not perpendicular to the streamlines, the slabs are immediately broken up, and the late stage coarsening (with a slowly-evolving, tubular structure along the streamlines) resembles that which we found above for the case of an initially homogenous phase, quenched to form demixing domains, in the presence of steady shear.

IV. CONCLUSIONS

We have described two problems, in the physics of immiscible binary fluids, where inertial effects are important. One is the late stages of demixing (in the absence of applied flow) where the internal dynamics of the system drives it to high Reynolds numbers. Whether the final Re is self-limiting, as recently suggested by Grant and Elder [4] remains to be seen; however, we have observed [5] clear evidence of an inertial-dominated regime in which the Furukawa ($l \sim t^{2/3}$) scaling is found. The second problem is in the steady state behaviour of binary fluids in simple shear. Our preliminary simulation results, though far from conclusive, inspired us to speculate that the very existence of such a steady state is itself dependent on inertial effects. If so, the scaling analysis that underlies Doi-Ohta theory [6], is in doubt. Future simulations on much larger systems should allow us to settle this point.

We thank Simon Jury, Patrick Warren and Julia Yeomans for valuable discussions, and Alistair Young for the turbulence dns simulation code. Work funded in part under the EPSRC E7 Grand Challenge.

-
- [1] A. J. Bray, *Adv. Phys.* **43**, 357 (1994).
 - [2] See, *e.g.* K. Kubota, N. Kuwahara, H. Eda and M. Sakazume, *Phys. Rev. A* **45**, R3377 (1992); S. H. Chen, D. Lombardo, F. Mallamace, N. Micali, S. Trusso, and C. Vasi, *Prog. Coll. Polymer Sci.* **93**, 331 (1993); T. Hashimoto, H. Jinnai, H. Hasegawa, and C. C. Han, *Physica A* **204**, 261 (1994).
 - [3] A. J. Wagner and J. M. Yeomans, *Phys. Rev. Lett.* **80**, 1429 (1998).
 - [4] M. Grant and K. R. Elder, *Phys. Rev. Lett.* **82**, 14 (1999).
 - [5] V. M. Kendon, J.-C. Desplat, P. Bladon and M. E. Cates *3D spinodal decomposition in the inertial regime*, *Phys. Rev. Lett.* **83**, 576 (1999), cond-mat/9902346
 - [6] M. Doi and T. Ohta, *J. Chem. Phys.* **95**, 1242 (1991).
 - [7] M. R. Swift, E. Orlandini, W. R. Osborn and J. Yeomans, *Phys. Rev. E* **54**, 5041 (1996); A. J. Wagner and J. M. Yeomans, *Phys. Rev. Lett.* **80**, 1429 (1998).
 - [8] R. D. Groot and P. B. Warren, *J. Chem. Phys.* **107**, 4423 (1997).
 - [9] P. Bladon and J.-C. Desplat, in preparation.
 - [10] S. I. Jury, P. Bladon, S. Krishna and M. E. Cates, *Phys. Rev. E*, **59**, R2535 (1999).
 - [11] E. D. Siggia, *Phys. Rev. A* **20**, 595 (1979).
 - [12] H. Furukawa, *Adv. Phys.* **34**, 703 (1985).
 - [13] M. Laradji, S. Toxvaerd, and O. G. Mouritsen, *Phys. Rev. Lett.* **77**, 2253 (1996).
 - [14] S. Bastea and J. L. Lebowitz, *Phys. Rev. Lett.* **78**, 3499 (1997).
 - [15] The linear law has been reported by a number of groups for which reliable parameter values are unavailable: T. Koga and K. Kawasaki, *Phys. Rev. A* **44**, R817 (1991); S. Puri and B. Dünweg, *Phys. Rev. A* **45**, R6977 (1992); F. J. Alexander, S. Chen, and D. W. Grunau, *Phys. Rev. B* **48**, 634 (1993); linear fits were not offered by W. Ma, A. Maritan, J. R. Banavar, J. Koplik, *Phys. Rev. A* **45**, R5347 (1992); A. Shinozaki and Y. Oono, *Phys. Rev. Lett.* **66**, 173 (1991), although we believe these data to be at low Re; C. Appert and S. Zaleski, *Phys. Rev. Lett.* **64**, 1 (1990) claimed to see the inertial region, but we doubt this; see [5].
 - [16] A. S. Monin and A. M. Yaglom, *Statistical Fluid Mechanics vol.2*, ed. J. Lumley, MIT Press, Cambridge MA (1975).
 - [17] V. M. Kendon, J.-C. Desplat, P. Bladon and M. E. Cates, in preparation.
 - [18] S. I. Jury, P. Bladon, M. E. Cates, S. Krishna, M. Hagen, N. Ruddock and P. B. Warren, *Phys. Chem. Chem. Phys.* *in press* (1999).
 - [19] R. G. Larson *The Structure and Rheology of Complex Fluids*, Oxford, NY (1999).
 - [20] Simulations in 3-D: T. Ohta, H. Nozaki, and M. Doi, *J. Chem. Phys.*, **93**, 2664, (1990); J. F. Olson and D. H. Rothman,

- J. Stat. Phys., **81**, 199, (1995); simulations in 2-D: P. Padilla and S. Toxvaerd, J. Chem. Phys., **106**, 2342, (1997); F. Corberi, G. Gonnella, and A. Lamura, Phys. Rev. Lett., **81**, 3852, (1998).
- [21] A. J. Wagner and J. M. Yeomans, *Phase Separation under Shear in Two-dimensional Binary Fluids*, to appear in Phys. Rev. E **59** (1999).
- [22] This is the main reason for the large wiggles; see also Y. Navot and M. Schwartz *Viscosity oscillations and hysteresis in dilute emulsions*, Phys. Rev. Lett. **79**, 4786 (1997) for a discussion of periodic breakup and reconnection of droplets.
- [23] B. J. Ackerson and N. A. Clarke, Phys. Rev. A **30**, 906 (1984).